

Michel Borghini and the rise of polarized targets


Colleagues and friends recall the work of a pioneer of a key technique in high-energy particle physics.

Michel Borghini, who passed away unexpectedly on 15 December 2012, was at CERN for more than 30 years. Born in 1934, Michel was a citizen of Monaco. He graduated from Ecole Polytechnique in 1955 and went on to obtain a degree in electrical engineering from Ecole Supérieure d'Electricité, Paris, in 1957. He then joined the group of Anatole Abragam at what was the Centre d'Etudes Nucléaires, Saclay, where he took part in the study of dynamic nuclear polarization that led to the development of the first polarized proton targets for use in high-energy physics experiments. It was here that he gained the experience that he was to develop at CERN, to the great benefit of experimental particle physics.

The basic aim with a polarized target is to line up the spins of the protons, say, in a given direction. In principle, this can be done by aligning the spins with a magnetic field but the magnitude of the proton's magnetic moment is such that it takes little energy to knock them out of alignment; thermal vibrations are sufficient. Even at low temperatures and reasonably high magnetic fields, the polarization achieved by this "brute force" method is small: only 0.1% at a temperature of 1 K and in an applied magnetic field of 1 T. To overcome this limitation, dynamic polarization exploits the much larger magnetic moment of electrons by harnessing the coupling of free proton spins in a material with nearby free electron spins. At temperatures of about 1 K, the electron spins are almost fully polarized in an external magnetic field of 1 T and the application of microwaves of around 70 GHz induces resonant transitions between the spin levels of coupled electron–proton pairs. The effect is to increase the natural, small proton polarization by more than two orders of magnitude. The polarization can be reversed with a slight change of the microwave frequency, with no need to reverse the external magnetic field.

First experiments

In 1962, Abragam's group, including Michel, reported on what was the first experiment to measure the scattering of polarized protons – in this case a 20 MeV beam derived from the cyclotron at Saclay – off a polarized proton target (Abragam *et al.* 1962). The target was a single crystal of lanthanum magnesium nitrate



Michel's 1968 paper on his "spin-temperature model".

($\text{La}_2\text{Mg}_3(\text{NO}_3)_{12} \cdot 24\text{H}_2\text{O}$ or LMN), with 0.2% of the La^{3+} replaced with Ce^{3+} , yielding a proton polarization of 20%.

Michel moved to CERN three years later, where he and others from Saclay and CERN had just tested a polarized target in an experiment on proton–proton scattering at 600 MeV at the Synchrocyclotron (SC) (CERN Courier December 2007 p12). Developed by the Saclay group for the higher energy beams of the Proton Synchrotron (PS), the target consisted of a crystal of LMN 4.5 cm long with transverse dimensions 1.2 cm $\times$ 1.2 cm and doped with 1% neodymium. It was cooled to around 1 K in a ^4He cryostat built in Saclay by Pierre Roubeau, in the field of a 1.8 T magnet designed by CERN's Guido Petrucci and built in the SC workshop. This target, with an average polarization of around 70%, was used in several experiments at the PS between 1965 and 1968, in both pion and proton beams with momenta of several GeV/c. These experiments measured the polarization parameter for π^+ elastic scattering and for the charge-exchange reaction $\pi^+ p \rightarrow \pi^0 n$ at small values of t , the square of the four-momentum transfer, typically, $|t| < 1 \text{ GeV}^2$.

In LMN crystals, the fraction of free, polarized protons is only around 1/16 of the total number of target protons. As a consequence, the unpolarized protons bound in the La, Mg, N and O nuclei formed a serious background in these early experiments. This background was reduced by imposing on the final-state particles the strict kinematic constraints expected from the collisions off protons at rest; the residual background was then subtracted by taking special data with a "dummy" target containing no free protons.

Michel's group at CERN thus began investigating the possibility of developing polarized targets with a higher content of free protons. In this context, in 1968 Michel published two important

Tribute

papers in which he proposed a new phenomenological model of dynamic nuclear polarization: the “spin-temperature model” (Borghini 1968a and 1968b). The model suggested that sizable proton polarizations could be reached in frozen organic liquids doped with paramagnetic radicals. Despite some initial scepticism, in 1969 Michel’s team succeeded in measuring a polarization of around 40% in a 5 cm³ sample consisting of tiny beads made from a frozen mixture of 95% butanol (C₄H₉OH) and 5% water saturated with the free-radical porphyrin. The beads were cooled to 1 K in an external magnetic field of 2.5 T and the fraction of free, polarized protons in the sample was around 1/4 – some four times higher than in LMN (Mango, Runólfsson and Borghini 1969).


The group at CERN went on to study a large number of organic materials doped with free-paramagnetic radicals, searching for the optimum combination for polarized targets. In this activity, where cryostats based on ³He–⁴He dilution capable of reaching temperatures below 0.1 K were developed, Michel guided two PhD students: Wim de Boer of the University of Delft (now professor at the Karlsruhe Institute of Technology) and Tapio Niinikoski of the University of Helsinki, who went on to join CERN in 1974. They finally obtained polarizations of almost 100% in samples of propanediol (C₃H₈O₂) doped with chromium (V) complexes and cooled to 0.1 K, in a field of 2.5 T, with 19% free, polarized protons.

In this work, the concept of spin temperature that Michel had proposed was verified by polarizing several nuclei simultaneously in a special sample containing ¹³C and deuterons. The nuclei had different polarizations but their values corresponded to a single spin temperature in the Boltzmann formula giving the populations of the various spin states.

These targets were used in a number of experiments at CERN, at both the PS and the Super Proton Synchrotron (SPS). They measured polarization parameters in the elastic scattering of pions, kaons and protons on protons in the forward diffraction region and at backward scattering angles; in the charge-exchange reactions $K^+p \rightarrow \bar{K}^0n$ and $\bar{p}p \rightarrow \bar{n}n$; in the reaction $\pi^+p \rightarrow K^0\Lambda^0$; and in proton–deuteron scattering. In all of these experiments, Jean-Michel Rieubland of CERN provided invaluable help to ensure a smooth operation of the targets.

In the early 1970s, Michel also initiated the development of “frozen spin” targets. In these targets, the proton spins were first dynamically polarized in a high, uniform magnetic field, and then cooled to a low enough temperature so that the spin-relaxation rate of the protons would be slow even in lower magnetic fields. The targets could then be moved to the detector, thus providing more freedom in the choice of magnetic spectrometers and orientations of the polarization vector. The first frozen spin target was successfully operated at CERN in 1974.

In 1969, Michel took leave from CERN to join the Berkeley group led by Owen Chamberlain working at SLAC, where he took part in a test of T-invariance in inelastic e⁺ scattering from polarized protons in collaboration with the SLAC group led by Richard Taylor. The target, built at Berkeley, was made of butanol and the SLAC 20 GeV spectrometer served as the electron (and positron) analyser. The experiment measured the up–down asymmetry for transverse target spin for both electrons and positrons. No time-reversal violations were seen at the few per cent level.



Michel with a frozen-spin target at CERN in 1976.



The target's internal structure.



Michel during his time with the CERN Staff Association.

Michel took leave to work at SLAC again in 1977, this time on a search for parity violation in deep-inelastic scattering of polarized electrons off an unpolarized deuterium target. Here, he worked on the polarized electron source and its associated laser, as well as on the electron spectrometer. The small parity-violation effects expected from the interference of the photon and Z exchanges were, indeed, observed and published in 1978. Michel then moved to the University of Michigan at Ann Arbor, where he joined the group led by Alan Krisch and took part in an experiment to measure proton–proton elastic scattering using both a polarized target and a 6 GeV polarized beam from the 12 GeV Zero Gradient Synchrotron at Argonne National Laboratory.

Michel left CERN’s polarized target group in 1978, succeeded by Niinikoski. Writing in 1985 on major contributions to spin physics, Chamberlain listed the people that he felt to be “the heroes – the people who have given [this] work a special push” (Chamberlain 1985). Michel is the only one that he cites twice: with Abragam and colleagues for the first polarized target and the first experiment to use such a target; and with Niinikoski, for their introduction of the

Tribute

frozen spin target and showing the advantages of powerful (dilution) refrigerators. Today, polarized targets with volumes of several litres and large ^3He – ^4He dilution cryostats are still in operation, for example in the NA58 (COMPASS) experiment at the SPS, where the spin structure of the proton has been studied using deep-inelastic scattering of high-energy muons (*CERN Courier* July/August 2006 p15 and September 2010 p34). Dynamic nuclear polarization has also found applications in medical magnetic-resonance imaging and Michel's spin-temperature model is still widely used.

In the 1980s, Michel took part in the UA2 experiment at CERN's SPS proton–antiproton collider, where he contributed to the calibration of the 1444 analogue-to-digital converters (ADCs) that were used to measure the energy deposited in the central calorimeter. He wrote all of the software to drive the large number of precision pulse-generators that monitored the ADC stability during data-taking.

From 1983 to 1996, he was a member of the Executive Committee of the CERN Staff Association, being its vice-president until 1990 and then its president until June 1996. After retiring from CERN in January 1999, he returned to Monaco where in 2003 he was nominated Permanent Representative of Monaco to the United Nations (New York), a post that he kept until 2005.

Michel was an outstanding physicist, equally at ease with theory and being in the laboratory. He had broad professional competences, a sharp, analytical mind, imagination and organizational skills. He is well remembered by his collaborators for his wisdom and advice, and also for his quiet demeanour and his keen but often subtle, sense of humour. His culture and interests extended well beyond science. He was also a talented tennis player. He will be sorely missed by those who had the privilege of working with him, or of being among his friends. Much sympathy goes to his two daughters, Anne and Isabelle, and to their families.

● Further reading

A Abragam *et al.* 1962 *Phys. Lett.* **2** 310.

M Borghini 1968a *Phys. Lett.* **26A** 242.

M Borghini 1968b *Phys. Rev. Lett.* **20** 419.

O Chamberlain 1985 *J. Phys. Colloques* **46 C2** 743.

S Mango, Ö Runólfsson and M Borghini 1969 *Nucl. Instr. Meth.* **72** 45.

Résumé

Michel Borghini et l'avènement des cibles polarisées

Michel Borghini, qui nous a quittés en décembre 2012, a été le pionnier d'une technique essentielle de la physique des particules. Il a commencé sa carrière au Centre d'études nucléaires de Saclay, où il a travaillé avec Anatole Abragam sur les études de polarisation nucléaire dynamique qui ont abouti aux premières cibles de protons polarisés pour la physique des hautes énergies. Au CERN, le groupe de Borghini a réalisé des avancées importantes dans le développement de cibles à base de liquides organiques gelés, pouvant permettre des polarisations de protons élevées. Dans cet article, ses collègues et amis rappellent différents aspects de ses contributions aux expériences, au CERN et ailleurs.

His colleagues and friends, at CERN and elsewhere.

Measuring magnetic field transients?

The Fast Digital Integrator FDI2056 is the world's fastest and most sensitive voltage integrator.

For the first time it is possible to measure fast, low-level magnetic field disturbances such as eddy currents in a switched magnet.

A complete and upgradable solution for your most exacting magnetic measurements.

Now it's possible!

Designed at the CERN



www.metrolab.com

METROLAB
Magnetic precision has a name

JANIS

Cryogenic Systems



Does your research require low temperatures? Contact Janis today. Our engineers will assist you in choosing the best system for your application.

- 10 mK to 800 K
- Cryocoolers
- LHe/LN₂ Cryostats
- Magnet Systems
- Dilution Refrigerator Systems
- Micro-manipulated Probe Stations

Janis Research Company

2 Jewel Drive Wilmington, MA 01887 USA

TEL +1 978 657-8750 FAX +1 978 658-0349 sales@janis.com

Visit our website at www.janis.com



Photomultipliers from ET Enterprises and ADIT Electron Tubes


Need to detect light down to single photon level?

Need dark counts as low as a few cps at 20° C?

Need a detection area of up to 200cm² or more?

If the answer is yes to any of these questions, then ET Enterprises, and sister company ADIT Electron Tubes in the US, may have the answer with their range of photomultipliers and associated electronics and hardware.

Photomultipliers have always offered significant performance advantages over other types of low-level light detection devices and this is as true today as ever before. But what has changed is that photomultipliers are now much easier to use due to the wide range of ready-to-use associated electronics and HV supplies, and modular analogue and photon counting systems, in many cases operating from low voltage.



Examples of how we can make using photomultipliers easier:

- ▮ Sockets and voltage dividers for almost any photomultiplier type and application.
- ▮ Magnetic shields.
- ▮ Modular signal processing electronics, analogue or digital.
- ▮ HV power supplies, including bench-top and compact, low power integral HV bases.
- ▮ Light-tight housings including cooling.

And, of course, we can also supply the photomultipliers for your application. We offer a very wide range, including alternatives to many Photonis types.

ET Enterprises Limited, Riverside Way,
Uxbridge, UB8 2YF, UK
Phone: +44 (0)1895 200880
Fax: +44 (0)1895 270873
sales@et-enterprises.com
www.et-enterprises.com

ADIT Electron Tubes, 300 Crane Street,
Sweetwater, Texas 79556, USA
Phone: (325) 235 1418
Fax: (325) 235 2872
sales@electrontubes.com
www.electrontubes.com

Contact us to learn more about these easy-to-use, highly sensitive light detectors, or try the parameterised search facility on our website.

Advertising feature

A long term partnership in the ultra-high vacuum technology between SAES Group and CERN



SAES Group has been pioneering the development and manufacturing of special getter materials in a variety of vacuum devices since more than 70 years. Thanks to a close technical cooperation, many SAES getters pumping solutions have been tested and used by CERN since more than 30 years, paving the way to a larger diffusion of getter technology to meet UHV-XHV challenges in particle accelerators. Today, SAES Group is supplying the new generation of getter vacuum pumps NEXTorr®, that combine high sorption speed for all the gasses in a compact volume:

the unique characteristics of NEXTorr® will enable improved vacuum performances in particles accelerators as well as reduced footprint and savings in the maintenance area.

SAES Group is also licensee of the not evaporable getter (NEG) film coating technology developed in the 90s' at CERN. SAES Group supplied NEG coated chambers to the most important particles accelerators in Europe, US and Asia. This technology nicely complements the getter pumps product line, thus allowing SAES Group to offer a complete portfolio of solutions for Particle Accelerators and High Energy machines.



saes
group

www.saesgroup.com
email: neg_technology@saes-group.com

- ✓ Pumping speed 3600 l/s (H2)
- ✓ Capacity 980 Torr/l (H2)
- ✓ CF 150 / CF 200